A Single Photon Can Simultaneously Excite Two or More Atoms

Luigi Garziano[1,2], Vincenzo Macrì[1], Roberto Stassi[1,2],
Omar Di Stefano[1], Franco Nori[2,3], and Salvatore Savasta[1,2]

[1]*Dipartimento di Fisica e di Scienze della Terra,*
Università di Messina, I-98166 Messina, Italy,
[2]*CEMS, RIKEN, Saitama 351-0198, Japan, and*
[3]*Physics Department, The University of Michigan,*
Ann Arbor, Michigan 48109-1040, USA

arXiv:1601.00886v2 [quant-ph] 26 Jan 2016

Abstract

We consider two separate atoms interacting with a single-mode optical resonator. When the frequency of the resonator field is twice the atomic transition frequency, we show that there exists a resonant coupling between *one* photon and *two* atoms, via intermediate virtual states connected by counter-rotating processes. If the resonator is prepared in its one-photon state, the photon can be jointly absorbed by the two atoms in their ground state which will both reach their excited state with probability close to one. Like ordinary quantum Rabi oscillations, this process is coherent and reversible, so that two atoms in their excited state will undergo a downward transition jointly emitting a single cavity photon. This joint absorption and emission processes can also occur with three atoms. The parameters used to investigate this process correspond to experimentally demonstrated values in circuit quantum electrodynamics systems.

PACS numbers: 42.50.Pq, 42.50.Ct, 85.25.Cp, 84.40.Az

Multiphoton excitation and emission processes were predicted in 1931 by Maria Göppert-Mayer in her doctoral dissertation on the theory of two-photon quantum transitions [1]. Two-photon absorption consists in the simultaneous absorption of two photons of identical or different frequencies by an atom or a molecule. Two-photon excitation is now a powerful spectroscopic and diagnostic tool [2, 3]. One may wonder if the reverse phenomenon, i.e., joint multiatom emission of one photon or multiatom excitation with a single photon, is ever possible. We show that these processes, not only can be enabled by the strong correlation between the states of the atoms and those of the field occurring in cavity quantum electrodynamics (QED) [4], but they can even take place with probability approaching one.

Cavity QED investigates the interaction of confined electromagnetic field modes with natural or artificial atoms under conditions where the quantum nature of light affects the system dynamics [5, 6]. A high degree of manipulation and control of quantum systems can be reached in the strong-coupling regime, where the atom-field coupling rate is dominant with respect to the loss and decoherence rates. This paves the way for many interesting physical applications [6–9]. Cavity QED is also very promising for the realization of quantum gates [10–12] and quantum networks for quantum computational tasks [13–15]. Many of the proposed concepts, pioneered with flying atoms, have been adapted and further developed using superconducting artificial atoms in the electromagnetic field of microwave resonators, giving rise to the rapidly growing field of circuit QED, which is very promising for future quantum technologies [8, 9, 12, 16–19]. In these systems, coupling rates between an individual qubit and a single electromagnetic mode of the order of 10% of the unperturbed frequency of the bare subsystems have been experimentally reached [20–23]. Such a coupling rate is significantly higher than that obtained using natural atoms. Such an ultrastrong coupling (USC) opens the door to the study of the physics of virtual processes which do not conserve the number of excitations governed by the counter-rotating terms in the interaction Hamiltonian [24–33]. Recently, it has been shown that these excitation-number-nonconserving processes enable higher order atom-field resonant transitions, making possible coherent and reversible multiphoton exchanges between the qubit and the resonator [34–36].

Here we examine a quantum system constituted by two two-level atoms coupled to a single-mode resonator in the regime where the field-atom detuning $\Delta = \omega_c - \omega_q$ is large (ω_c and ω_q are the resonance frequency of the cavity mode and the qubit transition frequency).

We investigate the situation where the two qubits are initially in their ground state and one photon is present in the resonator, corresponding to the initial state $|g, g, 1\rangle$. We find that, if $\omega_c \approx 2\omega_q$, a single cavity photon is able to excite simultaneously two independent atoms. During this process no parametric down-conversion, splitting the initial photon into observable pairs of photons at frequency $\omega_c/2$, occurs. The cavity photon is directly and jointly absorbed by the two atoms. As shown in Fig. 1, the initial state $|g, g, 1\rangle$ goes to virtual intermediate states that do not conserve the energy, but comes back to the real final state $|e, e, 0\rangle$ that does conserve energy. If $\omega_c \approx 3\omega_q$ the simultaneous excitation of three atoms: $|g, g, g, 1\rangle \rightarrow |e, e, e, 0\rangle$ is also possible. If the coupling is sufficiently strong, even a higher number of atoms can be excited with a single photon. Owing to optical selection rules, the two-atom process requires parity-symmetry breaking of the atomic potentials, which can be easily achieved in superconducting artificial atoms [34, 37, 38]. On the contrary, the three-atom process does not need broken symmetry.

The Hamiltonian describing the system consisting of a single cavity mode interacting with two or more identical qubits with possible symmetry-broken potentials is given by [20, 31]

$$\hat{H}_0 = \hat{H}_q + \hat{H}_c + \lambda \hat{X} \sum_i (\cos\theta\, \hat{\sigma}_x^{(i)} + \sin\theta\, \hat{\sigma}_z^{(i)}) \,, \tag{1}$$

where $\hat{H}_q = (\omega_q/2) \sum_i \hat{\sigma}_z^{(i)}$ and $\hat{H}_c = \omega_c \hat{a}^\dagger \hat{a}$, describe the qubit and cavity Hamiltonians in the absence of interaction, $\hat{X} = \hat{a} + \hat{a}^\dagger$, $\hat{\sigma}_x^{(i)}$ and $\hat{\sigma}_z^{(i)}$ are Pauli operators for the ith qubit, and λ is the coupling rate of each qubit to the cavity mode. For $\theta = 0$ parity is conserved. For flux qubits, this angle, as well as the transition frequency ω_q, can be continuously tuned by changing the external flux bias [20, 37]. For the sake of simplicity, Eq. (1) describes identical qubits, but this is not an essential point. In contrast to the Jaynes-Cummings (JC) model, the Hamiltonian in Eq. (1) explicitly contains counter-rotating terms of the form $\hat{\sigma}_+^{(i)} \hat{a}^\dagger$, $\hat{\sigma}_-^{(i)} \hat{a}$, $\hat{\sigma}_z^{(i)} \hat{a}^\dagger$, and $\hat{\sigma}_z^{(i)} \hat{a}$. The first (second) term creates (destroys) two excitations while the third (fourth) term creates (destroy) one excitation. The presence of counter-rotating terms in the interaction Hamiltonian enables four different paths which, starting from the initial state $|g, g, 1\rangle$, reach the final state $|e, e, 0\rangle$ (see Supplemental Fig. S1 [39]). Each path includes three virtual transitions involving out-of-resonance intermediate states. Fig. 1 only displays the process that gives the main contribution to the effective coupling between the bare states $|g, g, 1\rangle$ and $|e, e, 0\rangle$. Higher-order processes, depending on the atom-field interaction strength, can also contribute. By applying standard third-order perturbation

theory, we obtain the following effective coupling rate, $\Omega_{\rm eff}/\omega_{\rm q} \equiv (8/3)(\lambda/\omega_{\rm q})^3 \sin\theta\cos^2\theta$. The analytical derivation of the effective coupling rate as a function of $\lambda/\omega_{\rm q}$ is presented in the Supplemental Material [39]. Already at a coupling rate $\lambda/\omega_{\rm q} = 0.1$, an effective (two qubits)-(one photon) coupling rate $\Omega_{\rm eff}/\omega_{\rm q} \sim 10^{-3}$ can be obtained.



FIG. 1. (Color online) Sketch of the process giving the main contribution to the effective coupling between the bare states $|g,g,1\rangle$ and $|e,e,0\rangle$, via intermediate virtual transitions. Here, the excitation-number nonconserving processes are represented by arrowed dashed line. The transition matrix elements are also shown.

We diagonalize numerically the Hamiltonian in Eq. (1) for the case of two qubits and indicate the resulting energy eigenvalues and eigenstates as $\hbar\omega_i$ and $|i\rangle$ with $i = 0, 1, \ldots,$ choosing the labelling of the states such that $\omega_k > \omega_j$ for $k > j$. We use a normalized coupling rate $\lambda/\omega_{\rm q} = 0.1$ and an angle $\theta = \pi/6$. Figure 2a shows the frequency differences $\omega_{i,0} = \omega_i - \omega_0$ for the lowest energy states as a function of the resonator frequency. Starting from the lowest excited states of the spectrum, a large splitting anticrossing around $\omega_{\rm c}/\omega_{\rm q} = 1$ can be observed (see arrows in Fig. 2a). It corresponds to the standard vacuum Rabi splitting, which appears also when neglecting the counter-rotating terms. The straight line at $E/\omega_{\rm q} = 1$ corresponds to the dark antisymmetric state $(|g,e,0\rangle - |e,g,0\rangle)/\sqrt{2}$. Even larger splitting anticrossings around $\omega_{\rm c}/\omega_{\rm q} = 1$ can be observed at higher E values. These correspond to the second and third rung of the JC ladder. We are interested in the region around $\omega_{\rm c}/\omega_{\rm q} = 2$, where the levels 3 and 4 display an apparent crossing at $E/\omega_{\rm q} \approx 2$. Actually, what appears as a crossing on this scale, it turns out to be a splitting anticrossing on an enlarged view as in Fig. 2b. Observing that just outside this avoided-crossing region one level remains flat as a function of $\omega_{\rm c}$ with energy $\omega \approx 2\omega_{\rm q}$, while the other grows as $\omega_{\rm c}$, this splitting clearly originates from the hybridization of the states $|e,e,0\rangle$ and $|g,g,1\rangle$. The resulting states are well approximated by the states $(|e,e,0\rangle \pm |g,g,1\rangle)/\sqrt{2}$. This splitting is not present in the

rotating-wave approximation (RWA), where the coherent coupling between states with a different number of excitations is not allowed, nor does it occur in the absence of symmetry breaking ($\theta = 0$). The normalized splitting has a value $2\ \Omega_{\rm eff}/\omega_{\rm q} = 1.97 \times 10^{-3}$, which is in good agreement with 2×10^{-3}, obtained within perturbation theory. This observed hybridization opens the way to the observation of weird effects such as the simultaneous excitations of *two* qubits with only *one* cavity photon. Such a coupling between the states $|e, e, 0\rangle$ and $|g, g, 1\rangle$ can be analytically described by the effective interaction Hamiltonian $H_{\rm eff} = -\Omega_{\rm eff}(|e, e, 0\rangle\langle g, g, 1| + \text{H.c.})$.

A key theoretical issue of the USC regime is the distinction between bare (unobservable) excitations and physical particles that can be detected [28, 40]. For example, when the counter-rotating terms are taken into account, the mean photon number in the system ground state becomes different from zero: $\langle 0|\hat{a}^\dagger\hat{a}|0\rangle \neq 0$. However, these photons are actually virtual [40] because they do not correspond to real particles that can be detected in a photon-counting experiment. The same problem holds for the excited states. According to these analyses, the presence of an n-photon contribution in a specific eigenstate of the system does not imply that the system can emit n photons when prepared in this state.

In order to fully understand and characterize this anomalous avoided crossing not present in the RWA, a more quantitative analysis is required. In the following, we therefore calculate the output signals and correlations which can be measured in a photodetection experiment. We fix the cavity frequency at the value where the splitting between level 3 and 4 is minimum. Instead of starting from the ideal initial state $(|3\rangle - |4\rangle)/\sqrt{2} \approx |g, g, 1\rangle$, more realistically, we consider the system initially in its ground state $|0\rangle \approx |g, g, 0\rangle$ and study the direct excitation of the cavity by an electromagnetic Gaussian pulse with central frequency $\omega_{\rm d} = (\omega_{4,0} + \omega_{3,0})/2$. In this strongly-dispersive regime, the resonator displays very low anharmonicity, so that for a strong system excitation as that induced by a π-pulse, higher-energy states of the resonator (as the state $|8\rangle \simeq |g, g, 2\rangle$) can be resonantly populated. This problem can be avoided by feeding the system with a single photon input or by probing the system in the weak-excitation regime. However, in order to achieve a deterministic transition $|g, g, 1\rangle \rightarrow |e, e, 0\rangle$, a useful route involves introducing a Kerr nonlinearity into the resonator, able to activate a photon blockade. In circuit QED this can be realized by introducing some additional Josephson junction, or coupling the resonator with weakly-detuned artificial atoms [41]. This additional nonlinearity can be described by the Hamiltonian term



FIG. 2. (Color online) (a) Frequency differences $\omega_{i,0} = \omega_i - \omega_0$ for the lowest energy eigenstates of Hamiltonian (1) as a function of $\omega_{\rm c}/\omega_{\rm q}$. Here we consider a normalized coupling rate $\lambda/\omega_{\rm q} = 0.1$ between the resonator and each of the qubits. We used $\theta = \pi/6$. The black arrows indicate the ordinary vacuum splitting arising from the coupling between the states $|g,g,1\rangle$ and $(1/\sqrt{2})(|g,e,0\rangle + |e,g,0\rangle)$. (b) Enlarged view of the spectral region delimited by a square in panel (a). This shows an avoided-level crossing, demonstrating the coupling between the states $|g,g,1\rangle$ and $|e,e,0\rangle$ due to the presence of counter-rotating terms in the system Hamiltonian.

$\hat{H}_{\rm K} = \mu\,\hat{a}^{\dagger\,2}\,\hat{a}^2$. The driving Hamiltonian, describing the system excitation by a coherent electromagnetic pulse is $\hat{H}_{\rm d}(t) = \mathcal{E}(t)\cos(\omega t)\hat{X}$, where $\mathcal{E}(t) = A\exp\left[-(t-t_0)^2/(2\tau^2)\right]/(\tau\sqrt{2\pi})$. Here, A and τ are the amplitude and the standard deviation of the Gaussian pulse, respectively. A includes the factor $\sqrt{\kappa}$, where κ is the loss rate through the cavity port. The system is thus under the influence of the total Hamiltonian $\hat{H} = \hat{H}_0 + \hat{H}_{\rm K} + \hat{H}_{\rm d}(t)$.

The output photon flux emitted by a resonator can be expressed as $\Phi_{\rm out} = \kappa\langle\hat{X}^-\hat{X}^+\rangle$,

where $\hat{X}^+ = \sum_{j,k>j} X_{jk}|j\rangle\langle k|$ and $\hat{X}^- = (\hat{X}^+)^\dagger$, with $X_{jk} \equiv \langle j|(\hat{a}^\dagger + \hat{a})|k\rangle$, are the positive and negative frequency cavity-photon operators [30, 36]. Neglecting the counter-rotating terms, or in the limit of negligible coupling rates, they coincide with $\hat{a}$ and $\hat{a}^\dagger$, respectively. The signal directly emitted from the qubit is proportional to the qubit mean excitation number $\langle \hat{C}^- \hat{C}^+ \rangle$, where $\hat{C}^\pm$ are the qubit positive and negative frequency operators, defined as $\hat{C}^+ = \sum_{j,k>j} C_{jk}|j\rangle\langle k|$ and $\hat{C}^- = (\hat{C}^+)^\dagger$, with $C_{jk} \equiv \langle j|(\hat{\sigma}_- + \hat{\sigma}_+)|k\rangle$ [30, 36]. Neglecting the counter-rotating terms, or in the limit of negligible coupling rates, they coincide with $\hat{\sigma}_-$ and $\hat{\sigma}_+$, respectively. In circuit QED systems, this emission can be detected by coupling the qubit to an additional microwave antenna [8].

Thanks to the photon-blockade effect, induced by the Kerr interaction $\hat{H}_{\rm K}$, it is possible to resonantly excite the split states $|3\rangle$ and $|4\rangle$ with a π-pulse, so that after the pulse arrival the population is completely transferred from the ground state to only these two energy levels. We use a pulse width $\tau = 1/(4\,\omega_{43})$. Figure 3a displays the numerically-calculated dynamics of the photon number $\langle \hat{X}^- \hat{X}^+ \rangle$, of the mean excitation number $\langle \hat{C}_1^- \hat{C}_1^+ \rangle$ for qubit 1 (which, of course, coincides with that of qubit 2), and of the two-qubit correlation $G_{\rm q}^{(2)} \equiv \langle \hat{C}_1^- \hat{C}_2^- \hat{C}_2^+ \hat{C}_1^+ \rangle$. Vacuum Rabi oscillations showing the reversible excitation exchange between the qubits and the resonator are clearly visible. We observe that, after a half Rabi period, $\Omega_{\rm eff}\, t = \pi/2$, the excitation is fully transferred to the two qubits which reach an excitation probability approaching one. Hence, not only the multiatom absorption of a single photon is possible, but it can essentially be deterministic. We observe that the single-qubit excitation $\langle \hat{C}_i^- \hat{C}_i^+ \rangle$ and $G_{\rm q}^{(2)}$ almost coincide at any time. This almost-perfect two-qubit correlation is a clear signature of the joint excitation: if one qubit gets excited, the probability that also the other one is excited is very close to one. In summary, an electromagnetic pulse is able, thanks to the photon blockade effect, to generate a *single* cavity-photon, which then gets jointly absorbed by a *couple* of qubits. The resonant coupling can be stopped at this time, e.g., by changing the resonance frequency of the qubits. If not, the reverse process starts, where two qubits jointly emit a *single* photon: $|e, e, 0\rangle \rightarrow |g, g, 1\rangle$. We observe that $\langle \hat{X}^- \hat{X}^+ \rangle$ is not exactly zero at the photon minima. This occurs because the two-qubit excited state, owing to the same processes inducing its coupling with the one-photon state, acquires a dipole transition matrix element, so that this state is able to emit photons. We find that (not shown here) this effects increase when increasing the atom-field coupling strength λ. In order to exclude that this joint qubit excitation does not



FIG. 3. (Color online) (a) Time evolution of the cavity mean photon number $\langle \hat{X}^- \hat{X}^+ \rangle$ (dotted blue curve), qubit 1 mean excitation number $\langle \hat{C}_1^- \hat{C}_1^+ \rangle$ (continuous black curve), and the zero-delay two-qubit correlation function $G_{\rm q}^{(2)} = \langle \hat{C}_1^- \hat{C}_2^- \hat{C}_2^+ \hat{C}_1^+ \rangle$ (dashed red curve) after the arrival of a π-like Gaussian pulse initially exciting the resonator. After the arrival of the pulse, the system undergoes vacuum Rabi oscillations showing the reversible joint absorption and re-emission of one photon by two qubits. $\langle \hat{C}_1^- \hat{C}_1^+ \rangle$ and $G_{\rm q}^{(2)}(t)$ are almost coincident. This perfect two-qubit correlation is a signature that the two qubits are jointly excited. (b) Time evolution of the cavity mean photon number (dotted blue curve), the qubit mean excitation number, and the two-qubit correlation as in (a), but including the effect of cavity damping and atomic decay. The corresponding rates are $\kappa = \gamma = 4 \times 10^{-5} \omega_{\rm q}$.

occur via more conventional paths, involving the creation of photon pairs and/or a 1-qubit-1-photon excitation, we have also calculated the photonic second-order correlation function $G_{\rm c}^{(2)} \equiv \langle (\hat{X}^-)^2 (\hat{X}^+)^2 \rangle$ and the qubit-cavity correlation $G_{\rm qc}^{(2)} \equiv \langle \hat{C}_i^- \hat{X}^- \hat{X}^+ \hat{C}_i^+ \rangle$. We find that their value is more than two orders of magnitude lower than that of the two-qubit correlation $G_{\rm q}^{(2)}$.

Figure 3a has been obtained without including loss effects. The influence of cavity field

damping and atomic decay on the process can be studied by the master equation approach. We consider the system interacting with zero-temperature baths. By using the Born-Markov approximation without the post-trace RWA [35], the resulting master equation for the reduced density matrix of the system is

$$\dot{\hat{\rho}} = i[\hat{\rho}(t), \hat{H}] + \kappa \mathcal{D}\,[\hat{X}^{+}]\hat{\rho} + \gamma \sum_{i} \mathcal{D}[\hat{C}_{i}^{+}]\,\hat{\rho}\,, \tag{2}$$

where the superoperator $\mathcal{D}$ is defined as $\mathcal{D}[\hat{O}]\hat{\rho} = \frac{1}{2}(2\hat{O}\,\hat{\rho}\,\hat{O}^{\dagger} - \hat{\rho}\,\hat{O}^{\dagger}\,\hat{O} - \hat{O}^{\dagger}\,\hat{O}\,\hat{\rho})$. We use $\kappa = \gamma = 3 \times 10^{-5}\,\omega_{\rm q}$. Figure 3b shows how the cavity losses and the atomic decay affects the system dynamics. As expected, the vacuum Rabi oscillations undergo damping and, as expected, the two-qubit correlation is more fragile to losses. Finally, we have also considered the case when the two qubits display different coupling rates with the resonator field. We used $\lambda_1/\omega_{\rm q} = 8 \times 10^{-2}$ and $\lambda_2/\omega_{\rm q} = 1.2 \times 10^{-1}$. We found that also in this case $\langle \hat{C}_1^{-}\hat{C}_1^{+}\rangle = \langle \hat{C}_2^{-}\hat{C}_2^{+}\rangle \simeq G_{\rm q}^{(2)}$. This result *further confirms the simultaneous and joint nature of this multiatom process*.

The processes described here can be observed by placing two superconducting artificial atoms at opposite ends of a superconducting transmission line resonator [42]. These multi-atom excitation and emission processes can find useful applications for the development of novel quantum technologies. Conditional quantum-state transfer is a first possible application: the quantum information stored in one of the two qubits can be transferred to the resonator conditioned by the state of the second qubit. We also observe that the quantum Rabi oscillations displayed in Fig. 3 imply that a hybrid entangled GHZ state, $(|g,g,1\rangle + |e,e,0\rangle)/\sqrt{2}$, can be obtained by an elementary quantum Rabi process after a time $t = \pi/(4\Omega_{\rm eff})$. This state can be stored by just changing the transition frequency of one of the two qubits. Besides possible applications, the puzzling results presented here, showing that *one* photon can divide its energy into *two* spatially-separated atoms, and that vacuum fluctuations [43] can induce separate atoms to behave as a single quantum entity (as testified by the one-photon transition matrix element acquired by the transition $|g,g\rangle \rightarrow |e,e\rangle$), provide new insights into the quantum aspects of the interaction between light and matter.

We hope that this proposal for the simultaneous excitation of two or three atoms with a single photon, might be effective in producing the simultaneous excitation of two or three referees with a single manuscript.

This work is partially supported by the RIKEN iTHES Project, the MURI Center for Dynamic Magneto-Optics via the AFOSR award number FA9550-14-1-0040, the IMPACT program of JST, a Grant-in-Aid for Scientific Research (A), and from the MPNS COST Action MP1403 Nanoscale Quantum Optics.

Supplemental Material:

DERIVATION OF THE EFFECTIVE HAMILTONIAN

According to standard time-dependent perturbation theory, for a constant perturbation switched-on at $t = 0$, the resulting transition rate can be expressed as

$$W_{i\to f} = \frac{2\pi}{\hbar}|V_{fi}^{\text{eff}}|^2\,\delta(E_f - E_i)\,, \tag{3}$$

where i and f label the initial and final states with corresponding energies E_i and E_f, and $V_{f\,i}^{\text{eff}}$ describes the effective coupling strength connecting the initial and final states. In the framework of first-order perturbation theory, this effective coupling strength coincides with the matrix element of the generic perturbing interaction $\hat{V}$: $V_{fi}^{\text{eff}} = V_{fi} = \langle f|\hat{V}|i\rangle$. If i and f are coupled only via third-order perturbation theory, the resulting effective coupling strength is,

$$V_{fi}^{\text{eff}} = \sum_{m,n}\frac{V_{fn}V_{nm}V_{mi}}{(E_i - E_m)(E_i - E_n)}\,. \tag{4}$$

In the case when the states $|n\rangle$ and $|m\rangle$ are virtual intermediate states that do not conserve energy, the only effect of the perturbation is to couple, via these virtual intermediate states, the initial and final states. The same coupling can be described by the effective Hamiltonian

$$H_{\text{eff}} = V_{fi}^{\text{eff}}\,|f\rangle\langle i| + \text{H.c.}\,, \tag{5}$$

with V_{fi}^{eff} provided by Eq. (2).

We observe that, applying first-order perturbation theory by using this effective Hamiltonian, we obtain the same result of standard third-order perturbation theory with the real perturbation $\hat{V}$. Hence, Eq. (4) describes the effective coupling strength between the energy-degenerate states $|g, g, 1\rangle$ and $|e, e, 0\rangle$. We consider the case $\omega_{\text{c}} \approx 2\omega_{\text{q}}$ and a perturbation of

the form:

$$\hat{V} = \lambda \hat{X} \sum_i (\cos\theta\, \hat{\sigma}_x^{(i)} + \sin\theta\, \hat{\sigma}_z^{(i)})\,. \tag{6}$$

After carefully inspecting all the possible intermediate states, we find that only the four paths shown in Fig. S1 can connect the states $|g,g,1\rangle$ and $|e,e,0\rangle$. Applying Eq. (4), we obtain,

$$\Omega_{\rm eff} \equiv -V_{fi}^{\rm eff} = \frac{8}{3}\sin\theta\cos^2\theta\left(\frac{\lambda}{\omega_{\rm q}}\right)^3. \tag{7}$$

Figure S2 displays the comparison of the magnitudes of the effective Rabi splitting $2\Omega_{\rm eff}/\omega_{\rm q}$ between the states $|g,g,1\rangle$ and $|e,e,0\rangle$ obtained analytically [Eq. (5)] via third-order perturbation theory and by the numerical diagonalization of the Hamiltonian in Eq. (1) (in the main text), as a function of the normalized interaction strength $\lambda/\omega_{\rm q}$. The agreement is very good, also for coupling strengths λ beyond 10% of the qubit transition frequency $\omega_{\rm q}$. This result confirms the $(\lambda/\omega_{\rm q})^3$ proportionality of the effective (one-photon)-(two-atoms) coupling predicted by the above analysis.



FIG. 4. (Color online) Coupling between the bare states $|g,g,1\rangle$ and $|e,e,0\rangle$ via intermediate virtual transitions. Here, the excitation-number nonconserving processes are represented by the arrowed red dashed lines. The transition matrix elements are also shown (in blue).



FIG. 5. (Color online) Comparison between the numerically-calculated normalized Rabi splitting (points) (corresponding to twice the effective coupling between one cavity photon and two independent atoms) and the corresponding calculation using third-order perturbation theory (continuous red curve).

[1] M. Göppert-Mayer, "Über elementarakte mit zwei quantensprüngen," Annalen der Physik **401**, 273–294 (1931).

[2] W. Denk, J. H. Strickler, and W. W Webb, "Two-photon laser scanning fluorescence microscopy," Science **248**, 73–76 (1990).

[3] P. T. C. So, C. Y. Dong, B. R. Masters, and K. M. Berland, "Two-photon excitation fluorescence microscopy," Annual review of biomedical engineering **2**, 399–429 (2000).

[4] S. Haroche and J. M. Raimond, *Exploring the quantum: Atoms, Cavities and Photons* (Oxford Univ. Press, 2006).

[5] S. Haroche, "Nobel lecture: Controlling photons in a box and exploring the quantum to classical boundary," Rev. Mod. Phys. **85**, 1083 (2013).

[6] S. Haroche, M. Brune, and J. M. Raimond, "Atomic clocks for controlling light fields," Physics Today **66**, 27 (2013).

[7] S. Deleglise, I. Dotsenko, C. Sayrin, J. Bernu, M. Brune, J. M. Raimond, and S. Haroche, "Reconstruction of non-classical cavity field states with snapshots of their decoherence," Nature

455, 510–514 (2008).

[8] M. Hofheinz, H. Wang, M. Ansmann, R. C. Bialczak, E. Lucero, M. Neeley, A. D. O'Connell, D. Sank, J. Wenner, J. M. Martinis, and A. N. Cleland, "Synthesizing arbitrary quantum states in a superconducting resonator," Nature **459**, 546–549 (2009).

[9] B. Vlastakis, G. Kirchmair, Z. Leghtas, S. E. Nigg, L. Frunzio, S. M. Girvin, M. Mirrahimi, M. H. Devoret, and R. J. Schoelkopf, "Deterministically encoding quantum information using 100-photon Schrödinger cat states," Science **342**, 607–610 (2013).

[10] A. Rauschenbeutel, G. Nogues, S. Osnaghi, P. Bertet, M. Brune, J. M. Raimond, and S. Haroche, "Coherent operation of a tunable quantum phase gate in cavity QED," Phys. Rev. Lett. **83**, 5166 (1999).

[11] S.-B. Zheng and G.-C. Guo, "Efficient scheme for two-atom entanglement and quantum information processing in cavity QED," Phys. Rev. Lett. **85**, 2392 (2000).

[12] J. Q. You and F. Nori, "Superconducting circuits and quantum information," Phys. Today **58 (II)**, 42 (2005).

[13] D. Felinto, C.-W. Chou, J. Laurat, E. W. Schomburg, H. De Riedmatten, and H. J. Kimble, "Conditional control of the quantum states of remote atomic memories for quantum networking," Nature Physics **2**, 844–848 (2006).

[14] H. J. Kimble, "The quantum internet," Nature **453**, 1023–1030 (2008).

[15] S. Ritter, C. Nölleke, C. Hahn, A. Reiserer, A. Neuzner, M. Uphoff, M. Mücke, E. Figueroa, J. Bochmann, and G. Rempe, "An elementary quantum network of single atoms in optical cavities," Nature **484**, 195–200 (2012).

[16] J. Q. You and F. Nori, "Quantum information processing with superconducting qubits in a microwave field," Phys. Rev. B **68**, 064509 (2003).

[17] A. Blais, R. S. Huang, A. Wallraff, S. M. Girvin, and R. J. Schoelkopf, "Cavity quantum electrodynamics for superconducting electrical circuits: An architecture for quantum computation," Phys. Rev. A **69**, 062320 (2004).

[18] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R. S. Huang, J. Majer, S. Kumar, S. M. Girvin, and R. J. Schoelkopf, "Strong coupling of a single photon to a superconducting qubit using circuit quantum electrodynamics," Nature **431**, 162–167 (2004).

[19] I. Chiorescu, P. Bertet, K. Semba, Y. Nakamura, C.J.P.M. Harmans, and J. E. Mooij, "Coherent dynamics of a flux qubit coupled to a harmonic oscillator," Nature **431**, 159–162 (2004).

[20] T. Niemczyk, F. Deppe, H. Huebl, E. P. Menzel, F. Hocke, M. J. Schwarz, J. J. García-Ripoll, D. Zueco, T. Hümmer, E. Solano, A. Marx, and R. Gross, "Circuit quantum electrodynamics in the ultrastrong-coupling regime," Nature Phys. **6**, 772–776 (2010).

[21] P. Forn-Díaz, J. Lisenfeld, D. Marcos, J. J. García-Ripoll, E. Solano, C.J.P.M. Harmans, and J. E. Mooij, "Observation of the Bloch-Siegert shift in a qubit-oscillator system in the ultrastrong coupling regime," Phys. Rev. Lett. **105**, 237001 (2010).

[22] A. Fedorov, A. K. Feofanov, P. Macha, P. Forn-Díaz, C. J. P. M. Harmans, and J. E. Mooij, "Strong coupling of a quantum oscillator to a flux qubit at its symmetry point," Phys. Rev. Lett. **105**, 060503 (2010).

[23] P. Forn-Díaz, G. Romero, C. J. P. M. Harmans, E. Solano, and J. E. Mooij, "Broken selection rule in the quantum Rabi model," arXiv preprint arXiv:1510.03379 (2015).

[24] S. De Liberato, D. Gerace, I. Carusotto, and C. Ciuti, "Extracavity quantum vacuum radiation from a single qubit," Phys. Rev. A **80**, 053810 (2009).

[25] Q. Ai, Y. Li, H. Zheng, C. P. Sun, *et al.*, "Quantum anti-Zeno effect without rotating wave approximation," Phys. Rev. A **81**, 042116 (2010).

[26] X. Cao, J. Q. You, H. Zheng, A. G. Kofman, and F. Nori, "Dynamics and quantum Zeno effect for a qubit in either a low-or high-frequency bath beyond the rotating-wave approximation," Phys. Rev. A **82**, 022119 (2010).

[27] X. Cao, J. Q. You, H. Zheng, and F. Nori, "A qubit strongly coupled to a resonant cavity: asymmetry of the spontaneous emission spectrum beyond the rotating wave approximation," New J. Phys. **13**, 073002 (2011).

[28] R. Stassi, A. Ridolfo, O. Di Stefano, M. J. Hartmann, and S. Savasta, "Spontaneous conversion from virtual to real photons in the ultrastrong-coupling regime," Phys. Rev. Lett. **110**, 243601 (2013).

[29] A. Ridolfo, S. Savasta, and M. J. Hartmann, "Nonclassical radiation from thermal cavities in the ultrastrong coupling regime," Phys. Rev. Lett. **110**, 163601 (2013).

[30] L. Garziano, A. Ridolfo, R. Stassi, O. Di Stefano, and S. Savasta, "Switching on and off of ultrastrong light-matter interaction: Photon statistics of quantum vacuum radiation," Phys. Rev. A **88**, 063829 (2013).

[31] L. Garziano, R. Stassi, A. Ridolfo, O. Di Stefano, and S. Savasta, "Vacuum-induced symmetry breaking in a superconducting quantum circuit," Phys. Rev. A **90**, 043817 (2014).

[32] J.-F. Huang and C. K. Law, "Photon emission via vacuum-dressed intermediate states under ultrastrong coupling," Phys. Rev. A **89**, 033827 (2014).

[33] Y.-J. Zhao, Y.-L. Liu, Y.-X. Liu, and F. Nori, "Generating nonclassical photon states via longitudinal couplings between superconducting qubits and microwave fields," Phys. Rev. A **91**, 053820 (2015).

[34] G. Zhu, D. G. Ferguson, V. E. Manucharyan, and J. Koch, "Circuit QED with fluxonium qubits: Theory of the dispersive regime," Phys. Rev. B **87**, 024510 (2013).

[35] K. K. W. Ma and C. K. Law, "Three-photon resonance and adiabatic passage in the large-detuning Rabi model," Phys. Rev. A **92**, 023842 (2015).

[36] L. Garziano, R. Stassi, V. Macrì, A. F. Kockum, S. Savasta, and F. Nori, "Multiphoton quantum Rabi oscillations in ultrastrong cavity QED," Phys. Rev. A **92**, 063830 (2015).

[37] Y. X. Liu, J. Q. You, L. F. Wei, C. P. Sun, and F. Nori, "Optical selection rules and phase-dependent adiabatic state control in a superconducting quantum circuit," Phys. Rev. Lett. **95**, 087001 (2005).

[38] F. Deppe, M. Mariantoni, E. P. Menzel, A. Marx, S. Saito, K. Kakuyanagi, H. Tanaka, T. Meno, K. Semba, H. Takayanagi, E. Solano, and R. Gross, "Two-photon probe of the Jaynes Cummings model and controlled symmetry breaking in circuit QED," Nature Phys. **4**, 686 (2008).

[39] See Supplemental Material for more details.

[40] A. Ridolfo, M. Leib, S. Savasta, and M. J. Hartmann, "Photon blockade in the ultrastrong coupling regime," Phys. Rev. Lett. **109**, 193602 (2012).

[41] A. J. Hoffman, S. J. Srinivasan, S. Schmidt, L. Spietz, J. Aumentado, H. E. Türeci, and A. A. Houck, "Dispersive photon blockade in a superconducting circuit," Phys. Rev. Lett. **107**, 053602 (2011).

[42] J. Majer, J.M. Chow, J.M. Gambetta, J. Koch, B.R. Johnson, J.A. Schreier, L. Frunzio, D.I. Schuster, A.A. Houck, A. Wallraff, *et al.*, "Coupling superconducting qubits via a cavity bus," Nature **449**, 443 (2007).

[43] P. D. Nation, J. R. Johansson, M. P. Blencowe, and F. Nori, "Stimulating uncertainty: Amplifying the quantum vacuum with superconducting circuits," Rev. Mod. Phys. **84**, 1–24 (2012).